

The most general methodology to create a valid correlation matrix for risk management and option pricing purposes

Riccardo Rebonato

Peter Jäckel

Quantitative Research Centre of the NatWest Group

19th October 1999

1 Introduction and motivation

The problem of how to specify a correlation matrix occurs in several important areas of finance and of risk management. A few of the important applications are, for instance, the specification of a (possibly time-dependent) instantaneous correlation matrix in the context of the BGM interest-rate option models, stress-testing and scenario analysis for market risk management purposes, or the specification of a correlation matrix amongst a large number of obligors for credit-derivative pricing or credit risk management.

For those applications where the most important *desideratum* is the recovery of the real-world correlation matrix, the problem is in principle well defined and readily solvable by means of well-established statistical techniques. In practice, however, the estimation problems can be severe: a small number of outliers, for instance, can seriously “pollute” a sample; non-synchronous data can easily destroy or hide correlation patterns; and the discontinuities in the correlation surface amongst forward rates when moving from deposit rates to the future strip, and from the latter to the swap market are well known to practitioners. In all these cases, the user often resorts to best-fitting the “noisy” elements of the sample correlation matrix by means of a plausible parametric function. This if, for instance, the route taken by Rebonato [8] for his calibration of the BGM

model.

Similarly, when faced with the problem of pricing credit derivatives such as, for instance, n^{th} -to-default credit swaps, practitioners often posit or adjust (rather than simply estimate) a correlation matrix between intuitively understandable factors, such as country, industrial sector, *etc.*

Even when the correlation matrix econometrically estimated is thought to be reliable, a risk manager often wants to alter it in an *ad hoc* fashion, either as a stress test or in the context of scenario analysis. It is well known, for instance, that, in the event of equity market crashes, the correlation between different equity markets can dramatically increase (see Boyer *et al.* [1]). A risk manager in charge of a portfolio made up of, say, several national equity indices would greatly overestimate the degree of diversification in his portfolio in the event of a crash if he used the matrix estimated during “normal” periods.

Another important situation where straightforward econometric estimation of a correlation matrix is not possible is the case of legacy currencies (such as French Franc, the Lira, *etc.*, which have all been subsumed in the Euro). To tackle this problem both the option pricer and the risk manager have to invent, on the basis of financial intuition, a correlation matrix for a currency for which no history exists.

Finally, and perhaps most importantly, statistical estimation techniques based on the analysis of historical data are intrinsically ill suited to provide predictions of future quantities when a paradigm shift has taken place. Such a paradigm shift could be, for instance, the imposition of currency controls for an emerging market, the granting of independence of a central Bank, a financial crisis such as Russia’s default in 1998, *etc.* In all these cases, it is to some extent possible to glimpse, from the traded prices of options, forward-looking estimates of volatilities that can supplement or replace the intrinsically backward-looking econometric estimates. The information available in derivative products about the future correlation matrix is, however, far too convoluted¹ to be of practical use. Whenever one therefore believed that the past might not

¹The most promising area might appear to be the caplet/swaption markets. Rebonato [6, 7, 8] however shows

be a reliable guide to the future, one could not realistically hope to distil a market consensus about the future correlation from the options market, and one would have to resort to altering an estimated matrix on the basis of financial intuition.

What all these different situations (and many more) have in common is therefore the desirability of altering a given correlation matrix deemed, for any reason, inadequate or inappropriate or of creating *tout court* a new one. The task, unfortunately, is not as simple as it might seem, given the requirement of a correlation matrix to be not only real symmetric, but also positive-semidefinite. This requirement, it must be stressed, does not belong to the “nice-to-have” list, but is an absolute requirement: if it were not fulfilled, for instance, a risk manager would have no guarantee that his VaR calculation, carried out using a variance-covariance normal approximation according to

$$VaR = k \cdot \mathbf{a}^T \cdot \mathbf{C} \cdot \mathbf{a} \quad (1)$$

will yield a positive number. In equation (1) above, \mathbf{a} is the vector of portfolio sensitivities to the market factors whose covariance matrix is denoted by \mathbf{C} and the constant k depends on the required percentile.

The problem has been recognized in the literature and several papers have appeared on the subject (see, *e.g.* Kupiec [4], Finger [3], Brooks *et al.* [2]). The solutions proposed so far, however, have been only partial answers to the problem. The technique proposed by Finger, for instance, is designed to increase portions of the correlation matrix. Finger’s approach (also reported in Brooks *et al.* [2]) has the drawback that some portions of the correlation matrix are altered in the desired fashion, but, in order to retain positive-semidefiniteness, the slack is taken up by other portions of the matrix in an uncontrolled fashion. The problems with this approach is clearly presented by Brooks *et al.*:

that, in the absence of serial options, it is impossible to pin down uniquely from the prices of traded options both the time-dependent volatility of the forward rates and the correlation matrix amongst them. Furthermore, even if one assumed perfect knowledge of the instantaneous volatility functions for all the forward rates, swaption prices have very poor discriminatory power with respect to different correlation matrices, because they depend more strongly on the functional form of the instantaneous volatilities.

“...It [Finger’s] is not a simple procedure in practice, and represents the main draw-back in J.P. Morgan’s approach [*i.e.* RiskMetrics] ... Note also that the other correlations in the matrix (that we had no *a priori* view on) have also changed, but this is the price that must be paid to ensure a consistent revised correlations matrix.”

The shrinkage technique (Kupiec [4]) is different in spirit and definitely more general. It proceeds by iterating towards a feasible solution that modifies as much as possible (in a non-easily quantifiable sense) a pre-existing well defined positive semidefinite matrix towards a desired target correlation matrix. Despite its appeal and greater generality, its main drawbacks are that it can be rather time consuming (since it requires a full matrix orthogonalisation at each iteration); that it requires as a starting point and “anchor” a positive-semidefinite matrix to start with; and that, above all, there is no way of determining to what extent the resulting matrix is optimal in any easily quantifiable sense.

In order to obviate these shortcoming we present a method which:-

- i) is guaranteed to produce a positive-semidefinite matrix;
- ii) does not require a pre-existing acceptable matrix to start with;
- iii) is fast to implement even for large matrices;
- iv) allows the determination of a feasible matrix that most closely approximates a target real symmetric (but not positive-semidefinite) matrix in a well-defined and quantifiable sense.

As for the last property of the method here presented, the user is at liberty to specify a suitable metric under which the technique is guaranteed to yield the best possible answer.

Furthermore, we present a second, and even faster, method, which shares properties i), ii) and iii) above, but which is not guaranteed to enjoy property iv)². In all the empirical studies we have carried out, however, the results obtained using this second approach have been extremely close,

²More precisely, there probably exists a metric with respect to which this method does provide the optimal solution but we have not been able to identify it.

albeit not identical, to the ones obtained using the first technique. After presenting the main method, we therefore propose that the latter can be used either as an excellent fast approximation to the solution obtained using what is probably the most intuitive metric or as an initial guess for the general problem.

2 Hypersphere decomposition

The starting point are the well known results from linear algebra that every $n \times n$ matrix \mathbf{M} given by

$$\mathbf{M} = \mathbf{W}\mathbf{W}^T \quad (2)$$

for any $\mathbf{W} \in \mathbb{R}^{n \times n}$ is positive-semidefinite and that, conversely, every positive-semidefinite matrix $\mathbf{M} \in \mathbb{R}^{n \times n}$ can be decomposed as in equation (2).

The method we propose for the construction of a valid correlation matrix

$$\hat{\mathbf{C}} = \mathbf{B}\mathbf{B}^T \quad (3)$$

that best-matches a given, *not* positive-semidefinite, target matrix \mathbf{C} is to view the elements of the row vectors of matrix \mathbf{B} in equation (3) as coordinates lying on a unit hypersphere [7, 8, 9]. If we denote by b_{ij} the elements of the matrix \mathbf{B} , the key is to obtain the $n \times n$ coordinates b_{ij} from $n \times (n - 1)$ angular coordinates θ_{ij} according to

$$\begin{aligned} b_{ij} &= \cos \theta_{ij} \cdot \prod_{k=1}^{j-1} \sin \theta_{ik} \quad \text{for } j = 1 .. n - 1 \\ &\text{and} \\ b_{ij} &= \prod_{k=1}^{j-1} \sin \theta_{ik} \quad \text{for } j = n . \end{aligned} \quad (4)$$

For an arbitrary set of angles $\{\theta_{ij}\}$, a matrix $\hat{\mathbf{C}}$ formed from \mathbf{B} as in equation (3) satisfies all the given constraints required of a correlation matrix by construction. In particular, thanks to the trigonometric relationship (4) and to the requirement that the radius of the unit hypersphere should be equal to one, the main diagonal elements are guaranteed to be unity.

In general, matrix $\hat{\mathbf{C}}$ will bear no resemblance to the target matrix \mathbf{C} . However, after using the above transformation and after defining a suitable error measure ε in the resulting approximate correlation matrix $\hat{\mathbf{C}}$

$$\varepsilon = \left\| \mathbf{C} - \hat{\mathbf{C}} \right\| , \quad (5)$$

one can use an optimisation procedure over the angles θ_{ij} to find the best possible fit given the chosen error measure. Sensible choices for the error measure are:-

- The sum of squares of the elements of the difference matrix $(\mathbf{C} - \hat{\mathbf{C}})$,

$$\chi^2_{\text{Elements}} := \sum_{ij} (c_{ij} - \hat{c}_{ij})^2 . \quad (6)$$

Since both \mathbf{C} and $\hat{\mathbf{C}}$ have unit diagonal elements, this error norm is equal to twice the sum of squares of errors in the free correlation coefficients.

- The elementwise sum of squares of errors in the sorted sets of eigenvalues of \mathbf{C} and $\hat{\mathbf{C}}$,

$$\chi^2_{\text{Eigenvalues}} := \sum_i (\lambda_i - \hat{\lambda}_i)^2 . \quad (7)$$

Naturally, the above suggestions are only examples and various other choices are conceivable. If, in particular, a risk manager felt that certain portions of the target correlation matrix \mathbf{C} should be recovered with particularly high accuracy, then correspondingly large weights could be assigned to the relative elements $(c_{ij} - \hat{c}_{ij})^2$. In this context, the error norm given by equation (6), where every element has exactly the same weight, has been shown by Rebonato [7, 8, 9] to have desirable global features in so far as the calibration of the BGM model is concerned.

The fundamental benefits of this method are twofold: first, when the underlying space over which the optimisation is carried out is expressed in terms of angle vectors describing coordinates on a unit hypersphere, no constraints have to be satisfied. This can be of substantial benefit for the numerical fitting procedure. More importantly, despite the fact that the procedure is still iterative, unlike the shrinking method [4], it simply requires a matrix multiplication per iteration,

rather than a full matrix diagonalisation³. This can make a big difference, especially for large matrices. Furthermore, the approach presented in the next section requires no iterations and provides a solution very similar to the one obtained using error metric (6). It can therefore be used to provide the starting point for the search procedure, thereby further reducing the overall computational cost.

3 Spectral decomposition⁴

Given the right-hand-side eigensystem \mathbf{S} of the real and symmetric matrix \mathbf{C} and its associated set of eigenvalues $\{\lambda_i\}$ such that

$$\mathbf{C} \cdot \mathbf{S} = \Lambda \cdot \mathbf{S} \quad \text{with} \quad \Lambda = \text{diag}(\lambda_i) \quad , \quad (8)$$

define the non-zero elements of the diagonal matrix Λ' as

$$\Lambda' : \lambda'_i = \begin{cases} \lambda_i & : \lambda_i \geq 0 \\ 0 & : \lambda_i < 0 \end{cases} . \quad (9)$$

If the target matrix \mathbf{C} is not positive-semidefinite, it has at least one negative eigenvalue and at least one of the λ'_i will be zero.

Also, define the non-zero elements of the diagonal scaling matrix \mathbf{T} with respect to the eigensystem \mathbf{S} by

$$\mathbf{T} : t_i = \left[\sum_m s_{im}^2 \lambda'_m \right]^{-1} . \quad (10)$$

Now, let

$$\mathbf{B}' := \mathbf{S} \sqrt{\Lambda'} \quad (11)$$

³It should be noted that for most optimisation procedures, the function defining the error norm has to be evaluated many times per iteration since typically some kind of Hessian or Jacobian matrix is also needed. Thus, the reduction in computational effort at this level can be crucial.

⁴also known as principal component analysis

and

$$\mathbf{B} := \sqrt{\mathbf{T}}\mathbf{B}' = \sqrt{\mathbf{T}}\mathbf{S}\sqrt{\Lambda'} \quad . \quad (12)$$

By construction,

$$\hat{\mathbf{C}} := \mathbf{B}\mathbf{B}^T \quad (13)$$

is now both positive-semidefinite and has unit diagonal elements. A procedural description of the above method may clarify what actually has to be done:

- Calculate the eigenvalues λ_i and the right-hand-side eigenvectors \mathbf{s}_i of \mathbf{C} .
- Set all negative λ_i to zero.
- Multiply the vectors \mathbf{s}_i with their associated “corrected” eigenvalues λ'_i and arrange as the columns of \mathbf{B}' .
- Finally, \mathbf{B} results from \mathbf{B}' by normalising the *row vectors* of \mathbf{B}' to unit length.

By following this procedure we obtain an acceptable correlation matrix which is intuitively “similar” to the target one (the more so, the fewer the eigenvalues which have to be set to zero). The crucial point, however, is not so much the plausibility of the metric but the fact that empirically we have always observed the results obtained using equations (8) to (13) to be very similar to those from the angular method discussed in section 2. How close the results are in practice is shown in the next section. This is significant because one can use the result of the method described here either as an accurate approximation to the best (in a χ^2_{Elements} sense) solution, or as the starting point for the optimisation discussed in section 2, thereby substantially reducing the computational burden of the hypersphere decomposition approach.

4 Examples

A risk manager retrieves from the middle office's reporting system the following correlation matrix of three world equity indices:

$$\tilde{\mathbf{C}} = \begin{pmatrix} 1 & 0.9 & 0.7 \\ 0.9 & 1 & 0.4 \\ 0.7 & 0.4 & 1 \end{pmatrix} .$$

The eigenvalues of $\tilde{\mathbf{C}}$ are $\{ -2.35364, 0.616017, 0.0303474 \}$ and the correlation matrix can be split up as in

$$\tilde{\mathbf{C}} = \tilde{\mathbf{B}}\tilde{\mathbf{B}}^T$$

with

$$\tilde{\mathbf{B}} = \begin{pmatrix} 0.98742 & 0.08718 & -0.13192 \\ 0.88465 & 0.45536 & 0.10021 \\ 0.77203 & -0.63329 & 0.05389 \end{pmatrix} .$$

The risk manager is aware of the Value at Risk calculated under the assumption of this correlation between the three indices. In order to assess the change in Value at Risk resulting from a decrease in correlation between two of the three underlying variables, the risk manager wishes to adjust the matrix to

$$\mathbf{C} = \begin{pmatrix} 1 & 0.9 & 0.7 \\ 0.9 & 1 & 0.3 \\ 0.7 & 0.3 & 1 \end{pmatrix} .$$

Unfortunately, the eigenvalues of \mathbf{C}' are now $\{ -2.29673, 0.710625, -0.00735244 \}$, and despite its plausible appearance, matrix \mathbf{C}' is no longer an acceptable correlation matrix. This highlights how a minor change can lead to the violation of the requirement of positive-semidefiniteness of a correlation matrix. The system will now fail when trying to construct a split-up

matrix \mathbf{B} for the purpose of Monte Carlo simulations⁵ in order to calculate the Value at Risk under the new assumptions.

Using the method outlined in section 2 with the error measure chosen to be χ^2_{Elements} as given by equation (6), we can calculate

$$\hat{\mathbf{B}} = \begin{pmatrix} 0.99804 & 0.06265 & 0 \\ 0.86482 & 0.50209 & 0 \\ 0.74020 & -0.67239 & 0 \end{pmatrix}$$

with

$$\hat{\mathbf{C}} = \hat{\mathbf{B}} \hat{\mathbf{B}}^T = \begin{pmatrix} 1 & 0.89458 & 0.69662 \\ 0.89458 & 1 & 0.30254 \\ 0.69662 & 0.30254 & 1 \end{pmatrix}$$

and a total error of $\chi^2_{\text{Elements}} = 0.946 \cdot 10^{-4}$.

In comparison, the method outlined in section 3 above, yields

$$\hat{\mathbf{B}} = \begin{pmatrix} 0.99805 & 0.06238 & 0 \\ 0.86434 & 0.50292 & 0 \\ 0.73974 & -0.67290 & 0 \end{pmatrix}$$

to give us

$$\hat{\mathbf{C}} = \hat{\mathbf{B}} \hat{\mathbf{B}}^T = \begin{pmatrix} 1 & 0.89402 & 0.69632 \\ 0.89402 & 1 & 0.30100 \\ 0.69632 & 0.30100 & 1 \end{pmatrix} .$$

One can notice that not only the total error of $\chi^2_{\text{Elements}} = 1.0 \cdot 10^{-4}$ but also the individual elements are remarkably close to the values obtained by optimisation. Despite the fact that there is in general no guarantee that the results of the two methods are as close together as in this example, we have always found very good agreement between the two approaches. To show the point,

⁵The construction of correlated normal variates from a vector of uncorrelated normal variates \mathbf{z} is done by the transformation $\mathbf{x} = \mathbf{B} \cdot \mathbf{z}$ with $\mathbf{C} = \mathbf{B} \mathbf{B}^T$.

we present below the stress case of a real symmetric 12×12 matrix which was constructed by calculating the statistical covariance of 316 vectors of 12 uniform random numbers and shifting the digits of 12 randomly chosen nondiagonal elements (randomly) by one place to the left or right (whilst limiting them to be $\in [-1, 1]$). Therefore, we have the given target

$$\mathbf{C} = \begin{pmatrix} 1.000 & -0.022 & -0.081 & -0.845 & 0.779 & 0.066 & -0.051 & 0.003 & -0.086 & 0.041 & 0.048 & -0.067 \\ -0.022 & 1.000 & -0.140 & -0.077 & 0.114 & -0.096 & -0.033 & -0.083 & 0.028 & 0.045 & 0.060 & -1.485 \\ -0.081 & -0.140 & 1.000 & 0.095 & -0.014 & -0.046 & -0.010 & -0.021 & -0.049 & 0.002 & -0.130 & 0.007 \\ -0.845 & -0.077 & 0.095 & 1.000 & -0.067 & 0.063 & 0.056 & -0.059 & 0.067 & 0.040 & -0.068 & 0.027 \\ 0.779 & 0.114 & -0.014 & -0.067 & 1.000 & -0.074 & 0.042 & 0.004 & -0.046 & 0.024 & 0.043 & -0.110 \\ 0.066 & -0.096 & -0.046 & 0.063 & -0.074 & 1.000 & 0.141 & 0.027 & -0.023 & 0.056 & -0.043 & 0.077 \\ -0.051 & -0.033 & -0.010 & 0.056 & 0.042 & 0.141 & 1.000 & 0.318 & -0.027 & -0.045 & 0.026 & 0.020 \\ 0.003 & -0.083 & -0.021 & -0.059 & 0.004 & 0.027 & 0.318 & 1.000 & -0.065 & -0.058 & 0.010 & 0.003 \\ -0.086 & 0.028 & -0.049 & 0.067 & -0.046 & -0.023 & -0.027 & -0.065 & 1.000 & -0.002 & -0.096 & -0.018 \\ 0.041 & 0.045 & 0.002 & 0.040 & 0.024 & 0.056 & -0.045 & -0.058 & -0.002 & 1.000 & -0.170 & -1.072 \\ 0.048 & 0.060 & -0.130 & -0.068 & 0.043 & -0.043 & 0.026 & 0.010 & -0.096 & -0.170 & 1.000 & -0.035 \\ -0.067 & -1.485 & 0.007 & 0.027 & -0.110 & 0.077 & 0.020 & 0.003 & -0.018 & -1.072 & -0.035 & 1.000 \end{pmatrix}$$

with the accompanying set of eigenvalues

$$\{2.899, 2.193, 1.370, 1.256, 1.065, 1.014, 0.946, 0.820, 0.762, 0.632, -0.135, -0.821\} .$$

We obtain from optimisation over angular coordinates

$$\hat{\mathbf{B}} = \begin{pmatrix} -0.179 & -0.967 & 0.039 & 0.162 & 0.056 & -0.011 & -0.019 & 0.035 & -0.033 & -0.002 & 0.000 & 0.000 \\ -0.905 & 0.110 & -0.028 & -0.279 & -0.034 & -0.078 & -0.018 & 0.213 & 0.186 & 0.040 & 0.000 & 0.000 \\ 0.096 & 0.134 & 0.111 & 0.450 & -0.647 & -0.047 & -0.002 & 0.518 & -0.256 & 0.056 & 0.000 & 0.000 \\ 0.135 & 0.766 & -0.055 & 0.054 & -0.127 & -0.081 & -0.542 & -0.223 & 0.064 & 0.137 & 0.000 & 0.000 \\ -0.219 & -0.671 & -0.012 & 0.145 & -0.187 & -0.254 & -0.576 & -0.168 & 0.084 & 0.116 & 0.000 & 0.000 \\ 0.090 & 0.016 & -0.349 & 0.320 & 0.547 & 0.422 & -0.283 & 0.402 & 0.038 & 0.218 & 0.000 & 0.000 \\ 0.064 & 0.018 & -0.796 & 0.032 & -0.034 & -0.211 & -0.123 & 0.093 & -0.021 & -0.540 & 0.000 & 0.000 \\ 0.075 & -0.070 & -0.732 & -0.007 & -0.154 & -0.217 & 0.346 & -0.156 & -0.045 & 0.486 & 0.000 & 0.000 \\ -0.011 & 0.159 & 0.201 & 0.029 & 0.494 & -0.732 & -0.023 & 0.170 & -0.348 & 0.062 & 0.000 & 0.000 \\ -0.674 & 0.161 & -0.027 & 0.489 & 0.083 & 0.190 & 0.069 & -0.334 & -0.341 & -0.069 & 0.000 & 0.000 \\ -0.006 & -0.154 & -0.112 & -0.697 & -0.090 & 0.275 & -0.245 & 0.045 & -0.574 & 0.053 & 0.000 & 0.000 \\ 0.988 & -0.136 & 0.053 & -0.004 & 0.024 & -0.011 & -0.015 & -0.028 & 0.021 & -0.012 & 0.000 & 0.000 \end{pmatrix}$$

with $\chi^2_{\text{Elements}} = 1.167$.

In comparison, using the spectral splitting technique, we calculate

$$\hat{\mathbf{B}} = \begin{pmatrix} -0.199 & -0.963 & 0.038 & 0.160 & 0.056 & -0.011 & -0.019 & 0.035 & -0.032 & -0.002 & 0.000 & 0.000 \\ -0.854 & 0.121 & -0.028 & -0.326 & -0.036 & -0.098 & -0.027 & 0.266 & 0.253 & 0.048 & 0.000 & 0.000 \\ 0.090 & 0.131 & 0.111 & 0.452 & -0.646 & -0.047 & -0.001 & 0.522 & -0.252 & 0.055 & 0.000 & 0.000 \\ 0.146 & 0.755 & -0.054 & 0.056 & -0.127 & -0.084 & -0.553 & -0.217 & 0.070 & 0.144 & 0.000 & 0.000 \\ -0.223 & -0.663 & -0.013 & 0.143 & -0.191 & -0.256 & -0.581 & -0.177 & 0.083 & 0.115 & 0.000 & 0.000 \\ 0.087 & 0.010 & -0.349 & 0.317 & 0.547 & 0.422 & -0.284 & 0.404 & 0.038 & 0.218 & 0.000 & 0.000 \\ 0.068 & 0.016 & -0.795 & 0.033 & -0.033 & -0.210 & -0.122 & 0.092 & -0.021 & -0.541 & 0.000 & 0.000 \\ 0.071 & -0.069 & -0.733 & -0.007 & -0.155 & -0.217 & 0.346 & -0.155 & -0.044 & 0.486 & 0.000 & 0.000 \\ -0.011 & 0.157 & 0.201 & 0.031 & 0.494 & -0.731 & -0.022 & 0.170 & -0.349 & 0.062 & 0.000 & 0.000 \\ -0.643 & 0.170 & -0.025 & 0.502 & 0.090 & 0.194 & 0.070 & -0.359 & -0.344 & -0.080 & 0.000 & 0.000 \\ -0.012 & -0.149 & -0.112 & -0.690 & -0.092 & 0.277 & -0.242 & 0.038 & -0.584 & 0.050 & 0.000 & 0.000 \\ 0.983 & -0.163 & 0.066 & -0.004 & 0.030 & -0.014 & -0.019 & -0.036 & 0.027 & -0.016 & 0.000 & 0.000 \end{pmatrix}$$

with $\chi^2_{\text{Elements}} = 1.208$ which demonstrates the proximity of the results of the two methods even for a doubly non-positive-semidefinite target matrix.

5 Conclusion

We have presented two simple methods to produce a feasible (*i.e.* real, symmetric, and positive-semidefinite) correlation matrix when the econometric one is either noisy, unavailable, or inappropriate. The first method is to the knowledge of the authors more general than any of the

approaches which have been proposed in the literature, and computationally faster. It can actually produce the optimal feasible solution in a sense specified by the user. The second method is, in principle, not as general, but we show that i) it is extremely fast and ii) it produces results very close to those obtained using the general procedure. It can therefore be used in its own right, or as a starting point for the general optimisation procedure, thereby making the latter even faster.

References

- [1] B. H. Boyer, M. S. Gibson, and M. Loretan. Pitfalls in tests for changes in correlations. International Finance Discussion paper #597, December 1997.
- [2] C. Brooks, B. Scott-Quinn, and J. Whalmsey. Adjusting VaR models for the impact of the Euro. Working paper of the ISMA Centre, 1998.
- [3] C. Finger. A methodology for stress correlation. In *RiskMetrics Monitor*, pages 3–11. Fourth Quarter 1997.
- [4] P. H. Kupiec. Stress testing in a value at risk framework. *Journal of Derivatives*, pages 7–24, Fall 1998.
- [5] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery. *Numerical Recipes in C*. Cambridge University Press, 1992.
- [6] Riccardo Rebonato. *Interest rate option models*. Jon Wiley and Sons, 1998.
- [7] Riccardo Rebonato. Calibration of the BGM model. *Risk*, March:74–79, 1999.
- [8] Riccardo Rebonato. On the simultaneous calibration of multifactor lognormal interest rate models to Black volatilities and to the correlation matrix. *Journal of computational finance*, 1999.
- [9] Riccardo Rebonato. *Volatility and Correlation*. Jon Wiley and Sons, 1999.